VANDERBILT UNIVERSITY MATH 196 — EXAMPLES OF SECTIONS 4.1 AND 4.2

Question 1. Determine whether the vectors (5, -2, 4), (2, -3, 5), and (4, 5 - 7) are linearly independent or dependent.

Question 2. Consider the set V of all triples (x, y, z) such that x = 3. Is V a vector space?

Question 3. Find solution vectors \vec{u} and \vec{v} such that the solution space is the set of all linear combinations of the form $s\vec{u} + t\vec{v}$:

$$\begin{cases} x_1 & -4x_2 & -3x_3 & -7x_4 & = 0\\ 2x_1 & -x_2 & +x_3 & +7x_4 & = 0\\ x_1 & +2x_2 & +3x_3 & +11x_4 & = 0 \end{cases}$$

SOLUTIONS.

1. Denote the vectors by $\vec{u} = (5, -2, 4), \ \vec{v} = (2, -3, 5), \ \text{and} \ \vec{w} = (4, 5 - 7).$ Consider

$$a\vec{u} + b\vec{v} + c\vec{w} = \vec{0}.$$

Recall that the vectors are linearly independent if the only solution of the previous equation is a = b = c = 0, and linearly dependent otherwise. The equation can be written as

$$a\begin{bmatrix}5\\-2\\4\end{bmatrix}+b\begin{bmatrix}2\\-3\\5\end{bmatrix}+c\begin{bmatrix}4\\5\\-7\end{bmatrix}=\begin{bmatrix}0\\0\\0\end{bmatrix},$$

or in matrix form

$$\begin{bmatrix} 5 & 2 & 4 \\ -2 & -3 & 5 \\ 4 & 5 & -7 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The system will have a unique solution provided that the matrix of the system is invertible. But we readily check that

$$\det \begin{bmatrix} 5 & 2 & 4 \\ -2 & -3 & 5 \\ 4 & 5 & -7 \end{bmatrix} = 0,$$

which means that the matrix is not invertible, hence the system does not have a unique solution, and therefore the vectors are linearly dependent.

2. First notice that elements of V can be written as (3, y, z). In order for V to be a vector space, there must exist a zero element, i.e., an element $q = (q_1, q_2, q_3)$ such that $q \in V$ and q + u = u for every $u \in V$. But if $q \in V$ then it can be written as $q = (3, q_2, q_3)$, and it follows that

$$q + u = (3, q_2, q_3) + (3, u_2, u_3) = (6, q_2 + u_2, q_3 + u_3) \neq (3, u_2, u_3).$$

Therefore V it is not a vector space.

3. The augmented matrix of the system is

$$\begin{bmatrix} 1 & -4 & -3 & -7 & \vdots & 0 \\ 2 & -1 & 1 & 7 & \vdots & 0 \\ 1 & 2 & 3 & 11 & \vdots & 0 \end{bmatrix}$$

Applying Gauss-Jordan elimination we find

Therefore x_3 and x_4 are free variables. Denoting by $x_3 = s$, $x_4 = t$, we can then write

$$x_1 = -s - 5t,$$

$$x_2 = -s - 3t.$$

Therefore solutions $\vec{x} = (x_1, x_2, x_3, x_4)$ can be written as

$$\begin{bmatrix} x_1\\x_2\\x_3\\x_4 \end{bmatrix} = \begin{bmatrix} -s-5t\\-s-3t\\s\\t \end{bmatrix} = s \begin{bmatrix} -1\\-1\\1\\0 \end{bmatrix} + t \begin{bmatrix} -5\\-3\\0\\1 \end{bmatrix} = s\vec{u} + t\vec{v},$$

where

$$\vec{u} = \begin{bmatrix} -1\\ -1\\ 1\\ 0 \end{bmatrix}, \ \vec{v} = \begin{bmatrix} -5\\ -3\\ 0\\ 1 \end{bmatrix}.$$

URL: http://www.disconzi.net/Teaching/MAT196-Spring-15/MAT196-Spring-15.html