

VANDERBILT UNIVERSITY  
MATH 208 — ORDINARY DIFFERENTIAL EQUATIONS  
EXAMPLES OF SECTION 2.4.

**Question 1.** Solve the following IVP:

$$\begin{cases} (4xy + 6y^2)y' + 3x^2 + 2y^2 = 0, \\ y(0) = 2. \end{cases}$$

**SOLUTION.** Write the equation as

$$(4xy + 6y^2)dy + (3x^2 + 2y^2)dx = 0,$$

and let  $M(x, y) = 3x^2 + 2y^2$ , and  $N(x, y) = 4xy + 6y^2$ . Computing we find  $\frac{\partial M}{\partial y} = 4y$  and  $\frac{\partial N}{\partial x} = 4y$ , and therefore this is an exact equation since  $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$ . Put

$$F(x, y) = \int M(x, y) dx = \int (3x^2 + 2y^2) dx = x^3 + 2xy^2 + g(y).$$

Then

$$\frac{\partial F}{\partial y} = 4xy + g' = N = 4xy + 6y^2 \Rightarrow g'(y) = 6y^2.$$

Integrating,  $g(y) = 2y^3$  (we do not need to add a constant here). Hence the general solution is

$$F(x, y) = x^3 + 2xy^2 + 2y^3 = C.$$

Using  $y(0) = 2$  we find  $C = 16$ , so the solution to the IVP is

$$x^3 + 2xy^2 + 2y^3 - 16 = 0.$$

Notice that this is an implicit solution.

*URL:* <http://www.disconzi.net/Teaching/MAT208-Fall-14/MAT208-Fall-14.html>