

VANDERBILT UNIVERSITY  
MATH 208 — ORDINARY DIFFERENTIAL EQUATIONS  
PRACTICE TEST 1 SOLUTIONS.

**Question 1.** For each equation below, identify the unknown function, classify the equation as linear or non-linear, and state its order.

(a)  $y \frac{dy}{dx} + \frac{y}{x} = 0$ .

**Solution.** Unknown:  $y$ . Non-linear, first order.

(b)  $x'''' + \cos t x' = \sin t$ .

**Solution.** Unknown:  $x$ . Linear, fourth order.

(c)  $y''' = -\cos y y'$ .

**Solution.** Unknown:  $y$ . Non-linear, third order.

**Question 2.** Solve the following initial value problems.

(a)  $y' = \frac{y-1}{x+3}$ ,  $y(-1) = 0$ .

**Solution.** This equation is separable, so

$$\frac{dy}{y-1} = \frac{dx}{x+3} \Rightarrow \int \frac{dy}{y-1} = \int \frac{dx}{x+3},$$

from what we obtain

$$|y-1| = C|x+3|,$$

or yet

$$y = 1 + C(x+3).$$

Using the initial condition we find  $C = -\frac{1}{2}$ , thus

$$y = 1 - \frac{1}{2}(x+3).$$

(b)  $y' = e^{-x} - 4y$ ,  $y(0) = \frac{4}{3}$ .

**Solution.** This is a linear equation with  $p(x) = 4$  and  $q(x) = e^{-x}$ . Using the formula for first order linear equations, we find

$$e^{\int p(x) dx} = e^{4x},$$

so that

$$y = \frac{1}{3}e^{-x} + Ce^{-4x}.$$

The initial condition gives  $C = 1$ , so

$$y = \frac{1}{3}e^{-x} + e^{-4x}.$$

**Question 3.** Solve the following differential equations.

(a)  $y' = \frac{\cos y \cos x + 2x}{\sin y \sin x + 2y}.$

**Solution.** Write

$$(\cos y \cos x + 2x)dx - (\sin y \sin x + 2y)dy = 0.$$

We readily verify that this equation is exact, with  $M(x, y) = \cos y \cos x + 2x$  and  $N(x, y) = -(\sin y \sin x + 2y)$ . Then

$$F(x, y) = \int M(x, y) dx = \sin x \cos y + x^2 + g(y).$$

From

$$\frac{\partial F}{\partial y} = N,$$

we find

$$g'(y) = -2y,$$

hence  $g(y) = -y^2$ . The general solution is

$$F(x, y) = \sin x \cos y + x^2 - y^2 = C.$$

(b)  $y' = 2x^{-1}y + x^2 \cos x, \quad x > 0.$

**Solution.** This is a linear equation with  $p(x) = -\frac{2}{x}$ , and  $q(x) = x^2 \cos x$ . Using the formula for first order equations, we find

$$y = x^2 \sin x + Cx^2.$$

(c)  $x^2 y' = y - 1.$

**Solution.** This is a separable equation:

$$\frac{dy}{y-1} = \frac{dx}{x^2}.$$

Integrating, we get

$$\ln |y-1| = -\frac{1}{x} + C,$$

which leads to

$$y = Ce^{-x^{-1}} + 1.$$

The solution  $y = 1$  is included in the above family upon taking  $C = 0$ .

**Question 4.** Find the general solution of the given differential equation.

(a)  $y'' + 8y' - 14y = 0.$

**Solution.**  $\lambda^2 + 8\lambda - 14 = 0 \Rightarrow \lambda = -4 \pm \sqrt{30}$ .  $y = c_1 e^{(-4+\sqrt{30})t} + c_2 e^{(-4-\sqrt{30})t}.$

(b)  $y'' + 8y' - 9y = 0$ .

**Solution.**  $\lambda^2 + 8\lambda - 9 = 0 \Rightarrow \lambda = -9, \lambda = 1$ .  $y = c_1e^{-9t} + c_2e^t$ .

(c)  $t^2y'' + 5y = 0, t > 0$ .

**Solution.** Cauchy-Euler equation.  $\lambda^2 - \lambda + 5 = 0 \Rightarrow \lambda = \frac{1 \pm i\sqrt{19}}{2}$ .  $y = c_1t^{\frac{1}{2}} \cos(\frac{\sqrt{19}}{2} \ln t) + c_2t^{\frac{1}{2}} \sin(\frac{\sqrt{19}}{2} \ln t)$ .**Question 5.** Give the form of the particular solution for the given differential equations. You do not have to find the values of the constants of the particular solution.

(a)  $y'' + 2y' - 3y = \cos x$ .

**Solution.** The characteristic equation is

$$\lambda^2 + 2\lambda - 3 = (\lambda - 1)(\lambda + 3) = 0.$$

Hence  $y_1 = e^x$  and  $y_2 = e^{-3x}$  are linearly independent solutions of the associated homogeneous equation. Since these do not involve  $\cos x$ , we have

$$y_p = A \cos x + B \sin x.$$

(b)  $y'' + 4y = 8 \sin 2t$ .

**Solution.** The characteristic equation is

$$\lambda^2 + 4 = 0.$$

Hence  $y_1 = \cos 2t$  and  $y_2 = \sin 2t$  are linearly independent solutions of the associated homogeneous equation. Therefore we need to take  $s = 1$ , so

$$y_p = At \cos t + Bt \sin t.$$

(c)  $y'' - 2y' + y = e^t \cos t$ .

**Solution.** The characteristic equation is

$$\lambda^2 - 2\lambda + 1 = (\lambda - 1)^2 = 0.$$

Hence  $y_1 = e^t$  and  $y_2 = te^t$  are linearly independent solutions of the associated homogeneous equation. Thus,

$$y_p = (A \cos t + B \sin t)e^t.$$

(d)  $y'' - y' - 12y = 2t^6e^{-3t}$ .

**Solution.** The characteristic equation is

$$\lambda^2 - \lambda - 12 = (\lambda + 3)(\lambda - 4) = 0.$$

Hence  $y_1 = e^{-3t}$  and  $y_2 = e^{4t}$  are linearly independent solutions of the associated homogeneous equation. We need to take  $s = 1$ , thus

$$y_p = t(a_6t^6 + a_5t^5 + a_4t^4 + a_3t^3 + a_2t^2 + a_2t^1 + a_0)e^{-3t}.$$

**Question 6.** Verify that the given functions are two linearly independent solutions of the corresponding homogeneous equation. Then, find a particular solution solving the non-homogeneous problem.

(a)  $x^2y'' - 2y = 3x^2 - 1$ ,  $x > 0$ ,  $y_1 = x^2$ ,  $y_2 = x^{-1}$ .

(b)  $(1-x)y'' + xy' - y = \sin x$ ,  $0 < x < 1$ ,  $y_1 = e^x$ ,  $y_2 = x$ .

**Solution.** The verification is done by plugging in the given functions into the equation, while the particular solution is found with the formula

$$y_p(t) = -y_1(t) \int \frac{y_2(t)f(t)}{W(t)} dt + y_2(t) \int \frac{y_1(t)f(t)}{W(t)} dt,$$

The important thing to remember here is that in order to use the above formula we need the coefficient of  $y''$  to be equal to one. So, for example, in (b) we need to write


$$y'' + \frac{x}{1-x}y' - \frac{1}{1-x}y = \frac{\sin x}{1-x}.$$

Using the formula we find

$$y_p(t) = -e^x \int \frac{xe^{-x} \sin x}{(1-x)^2} dx + x \int \frac{\sin x}{(1-x)^2} dx.$$

**Question 7.** Match the direction fields with the given differential equations.

- (a)  $y' = -\frac{y}{x}$       (b)  $y' = \cos x$       (c)  $y' = y(1 - 0.5y)$       (d)  $y' = x$



**Solution.** (a)  $\rightarrow$  D, (b)  $\rightarrow$  A, (c)  $\rightarrow$  C, (d)  $\rightarrow$  B.

**Question 8.** Show that the problem

$$3y' - x^2 + xy^3 = 0, y(1) = 6,$$

has a unique solution defined in some neighborhood of  $x = 1$ .

**Solution.** Write

$$y' = f(x, y), y(1) = 6,$$

where  $f(x, y) = \frac{x^2 - xy^3}{3}$ . Since  $f(x, y)$  and  $\partial_y f(x, y) = -xy^2$  are continuous in the neighborhood of  $(1, 6)$ , the result follows from the existence and uniqueness theorem for first order equations.

URL: <http://www.disconzi.net/Teaching/MAT208-Fall-14/MAT208-Fall-14.html>