

VANDERBILT UNIVERSITY

MATH 2300 – MULTIVARIABLE CALCULUS

Examples of section 13.3

Question 1. Suppose you start at the point $(0, 0, 3)$ and move 5 units along the curve $x = 3 \sin t$, $y = 4t$, $z = 3 \cos t$ in the positive direction. Where are you now?

Solution 1. Write $\mathbf{r}(t) = \langle 3 \sin t, 4t, 3 \cos t \rangle$, so that $\mathbf{r}'(t) = \langle 3 \cos t, 4, -3 \sin t \rangle$ and $|\mathbf{r}'(t)| = 5$. The point $(0, 0, 3)$ corresponds to $t = 0$. Calculating the arc length:

$$s(t) = \int_0^t |\mathbf{r}'(\tau)| d\tau = 5t.$$

Since we moved five units along the curve, $s(t) = 5$, or $5t = 5 \Rightarrow t = 1$. Plugging $t = 1$ into $\mathbf{r}(t)$ gives $(3 \sin 1, 4, 3 \cos 1)$.

Question 2. Find the curvature of the ellipse $x = 3 \cos t$, $y = 4 \sin t$, $z = 0$ at the points $(3, 0, 0)$ and $(0, 4, 0)$.

Solution 2. Compute $\mathbf{r}'(t) = \langle -3 \sin t, 4 \cos t, 0 \rangle$ and $\mathbf{r}''(t) = \langle -3 \cos t, -4 \sin t, 0 \rangle$. Then

$$|\mathbf{r}'|^3 = (\sqrt{9 \sin^2 t + 4 \cos^2 t})^3,$$

and

$$\kappa(t) = \frac{|(-3 \sin t)(-4 \sin t) - 4 \cos t(-3 \cos t)|}{(\sqrt{9 \sin^2 t + 4 \cos^2 t})^3} = \frac{12}{(9 \sin^2 t + 4 \cos^2 t)^{\frac{3}{2}}}.$$

$(3, 0, 0)$ corresponds to $t = 0$, so $\kappa(0) = 3/16$; $(0, 4, 0)$ corresponds to $t = \pi/2$ so $\kappa(\pi/2) = 4/9$.