

VANDERBILT UNIVERSITY

MATH 3120 – INTRO DO PDES

*Some results on convergence of Fourier series*

Here we collect some useful results about convergence of Fourier series. Their proofs can be found in many textbooks, e.g., [1, 2].

We will denote by  $f(x^+)$  and  $f(x^-)$  the right and left values of a function  $f$  at  $x$ , defined by

$$f(x^+) = \lim_{h \rightarrow 0^+} f(x + h),$$

and

$$f(x^-) = \lim_{h \rightarrow 0^-} f(x + h).$$

If  $f$  is continuous at  $x$ , we have that  $f(x^+) = f(x^-)$ , but in general these values need not to be equal. For instance, let

$$f(x) = \begin{cases} -1, & x < 0, \\ 1, & x \geq 0. \end{cases}$$

Then  $f(0^+) = 1$  and  $f(0^-) = -1$ .

As done in class, the Fourier series of a function  $f$  will be written as

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right).$$

**Theorem 1.** *Let  $f$  be a piecewise  $C^1$  function defined on  $[-L, L]$ . Then, for any  $x \in (-L, L)$ ,*

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right) = \frac{1}{2} (f(x^+) + f(x^-))$$

*For  $x = \pm L$ , the series converges to  $\frac{1}{2}(f(-L^+) + f(L^-))$ .*

Thus, the Fourier series of  $f$  at  $x$  converges to  $f(x)$  if  $f$  is continuous at  $x$ .

Next, we consider differentiation and integration of Fourier series.

**Theorem 2.** *Let  $f$  be continuous on  $[-L, L]$ . Suppose that  $f(-L) = f(L)$ , and that  $f$  is piecewise  $C^2$ . Then, the Fourier series of  $f'$  can be obtained from that of  $f$  by differentiation term-by-term. I.e., if*

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right),$$

then

$$f'(x) = \sum_{n=1}^{\infty} \left( a_n \left( \cos \frac{n\pi x}{L} \right)' + b_n \left( \sin \frac{n\pi x}{L} \right)' \right),$$

whenever  $f'(x)$  equals its Fourier series. Equivalently,

$$f'(x) = \sum_{n=1}^{\infty} \left( -a_n \frac{n\pi}{L} \sin \frac{n\pi x}{L} + b_n \frac{n\pi}{L} \cos \frac{n\pi x}{L} \right).$$

The assumption that  $f(-L) = f(L)$  means that we could think of  $f$  as the restriction to  $[-L, L]$  of a continuous  $2L$ -periodic function.

Finally,

**Theorem 3.** *Let  $f$  be a piecewise continuous function on  $[-L, L]$  and assume that*

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right).$$

Then, for any  $x \in [-L, L]$ , we have

$$\int_{-L}^x f(s) ds = \int_{-L}^x \frac{a_0}{2} ds + \sum_{n=1}^{\infty} \left( a_n \int_{-L}^x \cos \frac{n\pi s}{L} ds + b_n \int_{-L}^x \sin \frac{n\pi s}{L} ds \right).$$

We now illustrate how one can use finitely many terms of the Fourier series to approximate a function. I.e., instead of taking an infinite sum in the Fourier series, we consider only the first  $N$  terms:

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^N \left( a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right).$$

Figure 1 shows the function  $|x|$  for  $-\pi \leq x \leq \pi$  and its Fourier series with  $N = 2$  and  $N = 4$  terms.




FIGURE 1. Graph of the function  $|x|$  and its corresponding Fourier series consisting only of the first two and four terms.

Figure 2 shows the function

$$f(x) = \begin{cases} -1, & -\pi \leq x < 0, \\ 1, & 0 \leq x \leq 0. \end{cases}$$

and its Fourier series with  $N = 6$  and  $N = 18$  terms.

Figure 3 shows the function  $\frac{x}{2}$  for  $-\pi \leq x \leq \pi$  and extended to a  $2\pi$ -periodic function, and its Fourier series with  $N = 4$  and  $N = 8$  terms.

In all these cases, the larger the  $N$ , the better the approximation, as expected.




FIGURE 2. Graph of the function  $f$  and its corresponding Fourier series consisting only of the first six and 18 terms.




FIGURE 3. Graph of the function  $\frac{x}{2}$  ( $2\pi$ -periodic), and its corresponding Fourier series consisting only of the first four and eight terms.

#### REFERENCES

- [1] T. Myint-U. *Partial Differential Equations of Mathematical Physics*. Elsevier Science Ltd, 1980.
- [2] W. A. Strauss. *Partial Differential Equations: An Introduction 2nd Edition*. Wiley, 2007.