

MAT 155B - FALL 12 - EXAMPLES SECTION 11.8

Question: Find the radius of convergence and interval of convergence of the series.

$$(a) \sum_{n=1}^{\infty} \frac{x^n}{\sqrt{n}} \quad (b) \sum_{n=1}^{\infty} (-1)^n \frac{n^2 x^n}{2^n} \quad (c) \sum_{n=1}^{\infty} \frac{(4x+1)^n}{n^2}$$

Solutions.

(a) Identify $c_n = \frac{1}{\sqrt{n}}$, $a = 0$ and $a_n = \frac{x^n}{\sqrt{n}}$. Use the ratio test

$$\left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{\frac{x^{n+1}}{\sqrt{n+1}}}{\frac{x^n}{\sqrt{n}}} \right| = \left| \frac{x^{n+1}}{\sqrt{n+1}} \frac{\sqrt{n}}{x^n} \right| = \frac{\sqrt{n}}{\sqrt{n+1}} |x|$$

So

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{\sqrt{n}}{\sqrt{n+1}} |x| = |x|$$

because $\lim_{n \rightarrow \infty} \frac{\sqrt{n}}{\sqrt{n+1}} = 1$. Now set $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$ to find

$$|x| < 1$$

Therefore the radius of convergence is $R = 1$. Since $a = 0$, the interval of radius 1 centered at $a = 0$ is $(-1, 1)$. To find the interval of convergence we need to plug at the endpoints. Plug $x = -1$ to find

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$$

This is an alternating series with $b_n = \frac{1}{\sqrt{n}}$. By the alternating series test, it converges.

Plugging $x = 1$ we find

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$$

This is a p -series with $p = \frac{1}{2}$. By the p -series test, it diverges. So the interval of convergence is $[-1, 1)$.

(b) Identify $c_n = \frac{(-1)^n n^2}{2^n}$, $a = 0$, $a_n = \frac{(-1)^n n^2}{2^n} x^n$. Use ratio test

$$\left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{\frac{(-1)^{n+1} (n+1)^2 x^{n+1}}{2^{n+1}}}{\frac{(-1)^n n^2 x^n}{2^n}} \right| = \left| \frac{(-1)^{n+1} (n+1)^2 x^{n+1}}{2^{n+1}} \frac{2^n}{(-1)^n n^2 x^n} \right| = \frac{1}{2} \frac{(n+1)^2}{n^2} |x|$$

So

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{1}{2} \frac{(n+1)^2}{n^2} |x| = \frac{1}{2} |x|$$

because $\lim_{n \rightarrow \infty} \frac{(n+1)^2}{n^2} = 1$. Now set $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$ to find

$$|x| < 2$$

Therefore the radius of convergence is $R = 2$. Since $a = 0$, the interval of radius 2 centered at $a = 0$ is $(-2, 2)$. To find the interval of convergence we need to plug at the endpoints. Plug $x = -2$ to find

$$\sum_{n=1}^{\infty} (-1)^n \frac{n^2 (-2)^n}{2^n} = \sum_{n=1}^{\infty} n^2$$

which diverges by the divergence test. Analogously plugging $x = 2$ we find

$$\sum_{n=1}^{\infty} (-1)^n \frac{n^2 (2)^n}{2^n} = \sum_{n=1}^{\infty} (-1)^n n^2$$

which also diverges by the divergence test. Hence the interval of convergence is $(-2, 2)$.

(c) First notice that we have to write the series as

$$\sum_{n=1}^{\infty} \frac{(4x+1)^n}{n^2} = \sum_{n=1}^{\infty} \frac{[4(x+\frac{1}{4})]^n}{n^2} = \sum_{n=1}^{\infty} \frac{4^n (x+\frac{1}{4})^n}{n^2}$$

since the term with power of x has to be of the form $(x-a)^n$, i.e., without a number multiplying the x . Now identify $c_n = \frac{4^n}{n^2}$, $a = -\frac{1}{4}$ (a is negative because $(x-a)^n = (x+\frac{1}{4})^n$), and $a_n = \frac{4^n (x+\frac{1}{4})^n}{n^2}$. The ratio test gives

$$\left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{\frac{4^{n+1} (x+\frac{1}{4})^{n+1}}{(n+1)^2}}{\frac{4^n (x+\frac{1}{4})^n}{n^2}} \right| = \left| \frac{4^{n+1} (x+\frac{1}{4})^{n+1}}{(n+1)^2} \frac{n^2}{4^n (x+\frac{1}{4})^n} \right| = 4 \frac{n^2}{(n+1)^2} \left| x + \frac{1}{4} \right|$$

So

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} 4 \frac{n^2}{(n+1)^2} \left| x + \frac{1}{4} \right| = 4 \left| x + \frac{1}{4} \right|$$

because $\lim_{n \rightarrow \infty} \frac{n^2}{(n+1)^2} = 1$. Now set $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$ to find

$$\left| x + \frac{1}{4} \right| < \frac{1}{4}$$

Therefore the radius of convergence is $R = \frac{1}{4}$. Since $a = -\frac{1}{4}$, the interval of radius $\frac{1}{4}$ centered at $a = -\frac{1}{4}$ is $(-\frac{1}{2}, 0)$. To find the interval of convergence we need to plug at the endpoints. Plug $x = -\frac{1}{2}$ to find

$$\sum_{n=1}^{\infty} \frac{4^n (-\frac{1}{2} + \frac{1}{4})^n}{n^2} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$$

which is absolutely convergent by comparing with a p -series with $p = 2$. Plugging $x = 0$ gives

$$\sum_{n=1}^{\infty} \frac{1}{n^2}$$

which converges by the p -series test. So the interval of convergence is $[-\frac{1}{2}, 0]$.