

VANDERBILT UNIVERSITY

MATH 3120 – INTRO DO PDES

HW 3

Question 1. Consider the Cauchy problem for Burger's equation:

$$\begin{aligned}u_t + uu_x &= 0, \\u(x, 0) &= h(x),\end{aligned}$$

for $(x, t) \in (-\infty, \infty) \times (0, \infty)$.

- (a) Find conditions on h that guarantee that no shock waves will form.
- (b) Derive a necessary condition for the formation of a shock wave.

Question 2. Consider the eikonal equation:

$$u_x^2 + u_y^2 = n^2, \tag{1}$$

where $n = n(x, y)$ is a given function. The eikonal equation [has important applications in optics](#).

The goal of this problem is to show how the method of characteristics can be used to solve the eikonal equation, which is a fully non-linear first order PDE.

Assume that an initial condition for (1) is given in the form of a parametrized curve $\Gamma(s) = (x_0(s), y_0(s), u_0(s))$.

- (a) Show that (1) is equivalent to $(u_x, u_y, n^2) \cdot (u_x, u_y, -1) = 0$ and interpret this geometrically.
- (b) Using (a), explain why it makes sense to consider the following system of characteristic equations for $x = x(t, s)$, $y = y(t, s)$, and $u = u(t, s)$ (recall the geometric meaning of the characteristic curves)

$$\dot{x} = u_x \tag{2a}$$

$$\dot{y} = u_y \tag{2b}$$

$$\dot{u} = n^2 \tag{2c}$$

- (c) From equations (2) and (1), derive

$$\ddot{x} = \frac{1}{2} \partial_x n^2 \tag{3a}$$

$$\ddot{y} = \frac{1}{2} \partial_y n^2 \tag{3b}$$

$$\dot{u} = n^2 \tag{3c}$$

- (d) Show that the solution to (1) is given by

$$u(x(t, s), y(t, s)) = u(x_0(s), y_0(s)) + \int_0^t (n(x(\tau, s), y(\tau, s)))^2 d\tau,$$

where $(x(\tau, s), y(\tau, s))$ is a solution to (3a)-(3b).

Question 3. Solve (1) when $n(x, y) = 1$ and with initial condition $u = 1$ on the curve $y = 2x$.

Question 4. Consider

$$\begin{aligned} u_{tt} - c^2 u_{xx} &= 0 \text{ in } (-\infty, \infty) \times (0, \infty), \\ u(x, 0) &= f(x), \\ u_t(x, 0) &= g(x), \end{aligned} \tag{4}$$

where $c = 3$ and

$$f(x) = g(x) = \begin{cases} 1, & |x| \leq 2 \\ 0, & |x| > 2. \end{cases}$$

(a) Without finding a general formula for u , find $u(0, 2)$.

(b) Without finding a general formula for u , compute

$$\lim_{t \rightarrow \infty} u(x, t).$$

(c) Solve (4).

(d) Is the solution you found classical? Explain.

Question 5. Consider the following problem for the wave equation on the half-line, i.e., for $x \geq 0$ rather than $-\infty < x < \infty$.

$$\begin{aligned} u_{tt} - 4u_{xx} &= 0 \text{ in } (0, \infty) \times (0, \infty), \\ u(x, 0) &= x^2 \text{ for } 0 \leq x < \infty, \\ u_t(x, 0) &= 6x \text{ for } 0 \leq x < \infty, \\ u(0, t) &= t^2 \text{ for } t > 0. \end{aligned} \tag{5}$$

(a) Notice that now we have the condition $u(0, t) = t^2$ for $t > 0$, which was absent when $-\infty < x < \infty$. Explain why such a condition was introduced.

(b) Solve (5).